


JUVÉDERM® Injectable Gel Fact Sheet

- JUVÉDERM® is the *first* smooth-consistency gel formulation
- JUVÉDERM® is the *only* hyaluronic acid (HA) filler FDA approved to last up to 1 year*

Lasts
up to
1 year*

JUVÉDERM® injectable gel is the first smooth-consistency gel



The smooth-consistency gel of JUVÉDERM® vs the granular-consistency gel of other HA dermal fillers: 2.4x magnification (≈ 0.05 mL).†

†The clinical significance of the difference has not been established.

Innovative HYLACROSS™ technology

JUVÉDERM® injectable gel is the first and only HA filler produced with innovative HYLACROSS™ technology, which combines a high concentration of HA with a high degree of cross-linking, creating a 3D-matrix structure. This robust network structure increases the resistance to enzymatic degradation.¹ HYLACROSS™ technology results in a smooth, natural look and feel for patients.

The JUVÉDERM® family of fillers

The **ONLY** family of HA fillers FDA approved to correct moderate to severe wrinkles and folds for up to 1 year*

JUVÉDERM® Ultra

- Optimally formulated and cross-linked for versatility in contouring and volumizing wrinkles and folds

JUVÉDERM® Ultra Plus

- Robust formulation in a 20% thicker gel¹ for volumizing and correcting deeper wrinkles and folds, such as nasolabial folds (NLFs)

Ask your Allergan representative for ordering information today, or call Allergan Customer Service at 1-800-377-7790.

For more information, visit www.juvederm.com.

Please see reverse side for important treatment considerations.

¹In the United States, JUVÉDERM® injectable gel is indicated for injection into the mid-to-deep dermis for correction of moderate to severe facial wrinkles and folds (such as nasolabial folds).



1. Data on file, Allergan, Inc.

A Brief Description of Relevant Indications for Use, Contraindications, Warnings, Precautions, and Adverse Events for JUVÉDERM® Injectable Gel

Indication: In the United States, JUVÉDERM® injectable gel is indicated for injection into the mid-to-deep dermis for correction of moderate to severe facial wrinkles and folds (such as nasolabial folds).

Contraindications: JUVÉDERM® injectable gel should not be used in patients who have severe allergies marked by a history of anaphylaxis or history or presence of multiple severe allergies. JUVÉDERM® injectable gel should not be used in patients with a history of allergies to Gram-positive bacterial proteins.

Warnings: JUVÉDERM® injectable gel should not be injected into blood vessels. If there is an active inflammatory process or infection at specific injection sites, treatment should be deferred until the underlying process is controlled.

Precautions: The safety of JUVÉDERM® injectable gel for use during pregnancy, in breastfeeding females, or in patients under 18 years has not been established. The safety and effectiveness of JUVÉDERM® injectable gel for the treatment of areas other than facial wrinkles and folds (such as lips) have not been established in controlled clinical studies. Patients who are using substances that can prolong bleeding, such as aspirin or ibuprofen, as with any injection, may experience increased bruising or bleeding at injection site. Patients should inform their physician before treatment if they are using these types of substances. As with all skin-injection procedures, there is a risk of infection. JUVÉDERM® injectable gel should be used with caution in patients on immunosuppressive therapy, or therapy used to decrease the body's immune response, as there may be an increased risk of infection. The safety of JUVÉDERM® injectable gel in patients with a history of excessive scarring (eg, hypertrophic scarring and keloid formations) and pigmentation disorders has not been studied. If laser treatment, chemical peel, or any other procedure based on active dermal response is considered after treatment with JUVÉDERM® injectable gel, or if JUVÉDERM® injectable gel is administered before the skin has healed completely after such a procedure, there is a possible risk of an inflammatory reaction at the treatment site.

Adverse events: The most commonly reported side effects are temporary injection-site redness, swelling, pain/tenderness, firmness, lumps/bumps, and bruising. Most side effects are mild or moderate in nature, and their duration is short lasting (7 days or less).

Important: For full safety information, please visit www.juvederm.com or call Allergan Product Support at 1-877-345-5372.

CAUTION: This device is restricted to sale by or on the order of a physician.